CULTIVATION RECOMMENDATION

Marfona.

Marfona is a fresh table variety with a yellow skin and light yellow flesh.



- **Rather firm cooking**
- **Uniform tuber shape and size**
- **Good virus resistances**

Characteristics

General characteristics

Breeder: Konst Research B.V.
Crossing: PRIMURA x KO 51-0123
Breedersright: Expired
Seedling: KO 66-0384

Plant and tuber characteristics

Maturity: second early / 6,5
Light sprout colour: red violet / RV
Flower colour: white / W
Number of berries: average / 6
Foliage - initial development: quite fast / 7
Foliage - final development: quite good / 7,5
Foliage - firmness: quite firm / 7
Foliage - selectability: average / 6
Skin colour: yellow / Y
Flesh colour: light yellow / 6
Skin finish: average / 6
Tuber shape: oval round / OR
Regularity of tuber shape: quite regular / 6
Tuberisation: quite low / 5
Tuber size: quite large / 7,5
Regularity of size: quite regular / 6,5
Dormancy: quite short / 5,5

Sensitivity and quality

Secondary growth: quite sensitive / 6,5
Black spot (internal bruising): quite sensitive / 6,5
Growth cracks: quite sensitive / 6
Harvest damage: quite sensitive / 6
Metribuzin - pre-emergence: slightly to not sensitive / 8
Bentazon: - / -
Ethylene: 5% fewer to 5% more tubers / 5
Underwater weight: 349
Dry matter content: 19,2%
Cooking type: rather firm / AB
Discolouration after cooking: quite sensitive / 6
Raw discolouration: quite sensitive / 6
Fries quality: unsuitable / 5
Dry matter distribution: - / -
Crisping quality: unsuitable / 4,5
TGA-level:
Carvone oil: 5% to 15% more tubers / 6
Dormancy: quite short / 5,5

Agrico has taken all reasonable care in producing this variety description. However, please note that any information may be subject to change and Agrico accepts no responsibility for errors or omissions. In no event shall Agrico be liable for damages arising out of or related to the information contained within this description.

AGRICO

CULTIVATION RECOMMENDATION

How to grow

Fertilization and seed treatment [1]

Nitrogen: standard advice - 10%.
Phosphate: standard advice - 10%.
Potassium: standard advice + 10%.

Seed treatment

Can be planted directly out of the cold store, preferably after a short heat shock.Desprouting more than once can have a negative impact on yield and size regularity.

Planting distance

28/35 mm: 22 cm (60.500 plants/ha)
35/55 mm: 27 cm (50.000 plants/ha)

Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

Fungicides

A strict preventive spraying schedule against late blight is necessary.

Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least two weeks after haulm killing before harvest.

Storage temperature - Storage time

Preferred storage temperature is 5 ºC. Suitable for medium to long term storage.



Say potato, say Agrico.

Resistances

Virus - Leaf roll: slightly susceptible / 6
Virus - Yn: slightly susceptible / 7
Virus - Yntn (tuber): slightly susceptible / 7
Spraing: susceptible / 5
Late blight - foliage: susceptible / 5
Late blight - tuber: slightly susceptible / 7
Fusarium: susceptible / 5,5
Common scab: very susceptible / 4,5
Powdery scab: slightly susceptible / 6
Virus - A: slightly to not susceptible / 8
Virus - X: slightly susceptible / 7
Black dot: susceptible / 5
Erwinia: slightly susceptible / 6
Wart disease F1: resistant / 10
Wart disease F6: - / -
Wart disease F18: - / -
Golden nematode Ro 1/4: - / -
Golden nematode Ro 2/3: - / -
White nematode Pa 2: - / -
White nematode Pa 3: - / -

Agrico • Duit 15 • 8305 BB Emmeloord • The Netherlands +31 (0)527 63 99 11 info@agrico.nl agricopotatoes.com