

# Lugano.



Processing

Lugano is an early maincrop yellow flesh processing variety with good resistances to potato cyst nematodes.

- ▶ **Large tubers**
- ▶ **Multiple resistances**
- ▶ **Good storability**



## characteristics

### General characteristics

Breeder: F.J. Vos

Crossing: AR 00-2211 x AR 99-0669

Breedersright: 31-12-2048

Seedling: VO 07-0289

### Plant and tuber characteristics

Maturity: early maincrop / 5,5

Light sprout colour: red / R

Flower colour: white / W

Number of berries: below average / 7

Foliage - initial development: quite fast / 7

Foliage - final development: good / 8

Foliage - firmness: quite firm / 7,5

Foliage - selectability: quite easy / 7

Skin colour: yellow / Y

Flesh colour: light yellow / 6,5

Skin finish: average / 6

Tuber shape: oval long / OL

Regularity of tuber shape: quite regular / 6

Tuberisation: quite low / 5

Tuber size: large / 8

Regularity of size: regular / 7

Dormancy: long / 8

### Sensitivity and quality

Secondary growth: quite sensitive / 6

Black spot (internal bruising): quite sensitive / 6

Growth cracks: quite sensitive / 6,5

Harvest damage: very sensitive / 4,5

Metribuzin - pre-emergence: slightly to not sensitive / 8

Metribuzin - post-emergence: slightly sensitive / 7,5

Bentazon: slightly to not sensitive / 8

Ethylene: 5% to 15% more tubers / 6

Underwater weight: 394

Dry matter content: 21,4%

Cooking type: rather floury / B

Discolouration after cooking: quite sensitive / 6,5

Raw discolouration: quite sensitive / 6

Fries quality: good / 7

Dry matter distribution: good / 7,5

Crisping quality: below average / 5,5

TGA-level: 6,4

Carvone oil: 25% more tubers / 8

Dormancy: long / 8

# How to grow



## Fertilization and seed treatment <sup>1</sup>

**Nitrogen:** standard advice.

**Phosphate:** standard advice.

**Potassium:** standard advice + 20%.

## Seed treatment

Can be planted directly out of the cold store, preferably after a short heat shock. Desprouting more than twice can have a negative impact on yield and size regularity.

## Planting distance

28/35 mm: 22 cm (60.000 plants/ha)

35/55 mm: 30 cm (45.000 plants/ha)

50/55 mm: 36 cm (37.000 plants/ha)

## Weeds

A standard use of metribuzin (Sencor), before or after emerge, doesn't give foliage reaction or yield reduction.

## Fungicides

A preventive spraying schedule against late blight is advised.

## Harvest

Due to its sensitivity scores for black spot and/or harvest damage, wait at least three weeks after haulm killing before harvest.

## Storage temperature - Storage time

Preferred storage temperature is 5 °C. For processing purposes, do not store below 7 °C. Suitable for long term storage.



## Resistances

**Virus - Leaf roll:** - / -

**Virus - Yn:** susceptible / 5,5

**Virus - Yntn (tuber):** slightly susceptible / 7,5

**Spraing:** susceptible / 5,5

**Late blight - foliage:** susceptible / 5

**Late blight - tuber:** susceptible / 5,5

**Fusarium:** slightly susceptible / 6,5

**Common scab:** susceptible / 5,5

**Powdery scab:** slightly susceptible / 7,5

**Virus - A:** - / -

Say potato,  
say **Agrico.**

**Virus - X:** - / -

**Black dot:** slightly susceptible / 6

**Erwinia:** - / -

**Wart disease F1:** resistant / 10

**Wart disease F6:** slightly susceptible / 9

**Wart disease F18:** - / -

**Golden nematode Ro 1/4:** resistant / 9

**Golden nematode Ro 2/3:** resistant / 9

**Golden nematode Ro 5:** resistant / 9

**White nematode Pa 2:** resistant / 9

**White nematode Pa 3:** resistant / 9